

DISINFECT SURFACES

Gently and Effectively



SaniZide Plus® is a convenient, fast-acting, multi-purpose, broad-spectrum cleaner, disinfectant, and deodorizer for environmental surfaces. Our alcohol-free formulation is a non-flammable, non-corrosive, EPA registered, quaternary ammonium compound that meets non-hazardous shipping requirements. SaniZide Plus® controls mold and mildew, helps prevent cross-contamination, and helps you comply with the OSHA Bloodborne Pathogens Standard, which requires the use of an intermediate level (tuberculocidal) disinfectant where gross contamination is known to have occurred.

Item#	Description	Case Qty.
34800	SaniZide Plus® 4oz. spray bottle	24/case
34805	SaniZide Plus® 16oz. spray bottle	12/case
34810	SaniZide Plus® 32oz. spray bottle	6/case
34815	SaniZide Plus® 1 gal. refill bottle	4/case



Effective against MRSA, VRE, Tuberculosis, Influenza, Poliovirus, and Norovirus

Ideal for use in:

- Hospitals
- Medical or Dental Offices
- Ambulances
- Ambulatory Surgical Centers
- · Nursing Homes
- Laboratories
- Pharmacies
- Other Healthcare Related Institutions





SaniZide Plus® Efficacy Sheet

COVID-19 Information:

SaniZide Plus® (EPA Reg # 1839-83-67161) has demonstrated effectiveness against viruses similar to 2019 novel coronavirus (SARS-CoV-2) on hard non-porous surfaces. Therefore, this product can be used against 2019 novel coronavirus (SARS-CoV-2) when used in accordance with the directions for use against Human Rotavirus or Rhinovirus 39 on hard, non-porous surfaces.

Refer to the CDC website (https://www.cdc.gov/coronavirus/2019-ncov/index.html) for additional information.

Virucidal Activity

Test Methods:

- * U.S. E.P.A. Pesticide Assessment Guidelines, Subdivision G: Product Performance, Section 91-2(f), and Section 91-30 (d), (e), November, 1982.
- † Protocols for Testing the Efficacy of Disinfectants against Hepatitis B Virus (HBV) (EPA, Federal Register, Vol. 65, No. 166, 8/25/2000, p. 51828).
- ‡ Protocol for Testing Disinfectants against Hepatitis C Virus using Bovine Viral Diarrhea Virus as approved by the U.S. EPA on August 15, 2002.
- **Modified U.S. E.P.A. Pesticide Assessment Guidelines, Subdivision G: Product Performance, Section 91-2(f), and Section 91-30 (d), (e), November, 1982.

^{***}Per E.P.A. Guidance Document dated 10/31/09.

Test Organism	Contact Time
*Hepatitis A Virus (HAV)	10 minutes
†Hepatitis B Virus (HBV)	5 minutes
‡Hepatitis C Virus (HCV)	5 minutes
*HIV-1 (associated with AIDS)	1 minute
*Human Coronavirus	2 minutes
**Norovirus (Norwalk Virus)	30 seconds
*Poliovirus Type 1, strain Brunhilde	10 minutes
*SARS Associated Coronavirus (ZeptoMetrix)	2 minutes
*Avian Flu A (Strains H9N2 and H3N2)	2 minutes
***2009 H1N1 Influenza A Virus	2 minutes

Tuberculocidal Data

Test Method: AOAC Confirmative In Vitro Test for Determining Tuberculocidal Activity	
Test Organism	Contact Time
Mycobacterium boyis BCG (Organon Teknika)	5 minutes

Fungicidal Data

Test Method: AOAC

rest Organism	Contact Time
Trichophyton mentagrophytes	10 minutes

Bactericidal Data

Test Method: AOAC

Test Organism	Contact Time
Staphylococcus aureus	3 minutes
Salmonella (choleraesuis) enterica (ATC 10708)	3 minutes
Pseudomonas aeruginosa	
Escherichia coli (E. coli)	3 minutes
Salmonella (typhi) enterica (ATCC 6539)	3 minutes
Streptococcus pyogenes	3 minutes
Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA)	3 minutes
Methicillin resistant Staphylococcus aureus (MRSA)	3 minutes
Methicillin resistant Staphylococcus epidermidis (MRSE)	
Vancomycin resistant Enterococcus faecalis (VRE)	
Vancomycin intermediate resistant Staphylococcus aureus (VISA)	

For use on:

Floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, shower stalls, bathtubs, cabinets, and all hard non-porous inanimate surfaces normally treated with a disinfectant.

^{*}For plastic and painted surfaces, spot test on an inconspicuous area before use.